

## Gulf of Mexico Harmful Algal Bloom Bulletin

Region: Texas


Monday, 08 February 2016

NOAA National Ocean Service

NOAA Satellite and Information Service

NOAA National Weather Service

Last bulletin: Monday, February 1, 2016



Satellite chlorophyll image with possible *K. brevis* HAB areas shown by red polygon(s), when applicable. Points represent cell concentration sampling data from January 29 to February 5: red (high), orange (medium), yellow (low b), brown (low a), blue (very low b), purple (very low a), pink (present), and green (not present). Cell count data are provided by Texas Parks and Wildlife Department. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide:

http://tidesandcurrents.noaa.gov/hab/hab\_publication/habfs\_bulletin\_guide.pdf

Detailed sample information can be obtained through the Texas Parks and Wildlife Department at: http://www.tpwd.state.tx.us./landwater/water/environconcerns/hab/redtide/status.phtml

## **Conditions Report**

*Karenia brevis* (commonly known as Texas red tide) ranges from not present to background concentrations along the coast of Texas. No respiratory irritation is expected Monday, February 8 through Tuesday, February 16.

Check http://tidesandcurrents.noaa.gov/hab/beach\_conditions.html for recent, local observations.

## **Analysis**


\*\*Due to the upcoming federal holiday, the next bulletin will be issued on Tuesday, February 16.\*\*

Sampling from the Texas A&M University's Imaging FlowCytobot, located on the Port Aransas ship channel, indicates that *Karenia brevis* ranges between 'not present' and 'background' concentrations (TAMU; 2/1-2/8). For information on area shellfish restrictions, contact the Texas Department of State Health Services.

In recent MODIS Aqua imagery (2/6, shown left), elevated to high levels of chlorophyll (3 to 15  $\mu$ g/L) are present along the entire Texas coastline, with the lowest levels observed near the Texas/Mexico border increasing northeast along the coast up to Galveston. Elevated chlorophyll is not indicative of the presence of *K. brevis* and is most likely due to the resuspension of benthic chlorophyll and sediments along the coast.

Forecast models based on predicted near-surface currents indicate a potential maximum transport of 10 km north from the Port Aransas region from February 6 to 11.

Urízar, Keeney




Wind speed and direction are averaged over 12 hours from buoy measurements. Length of line indicates speed; angle indicates direction. Red indicates that the wind direction favors upwelling near the coast. Values to the left of the dotted vertical line are measured values; values to the right are forecasts. Wind observation and forecast data provided by NOAA's National Weather Service (NWS).

## Wind Analysis

**Port Aransas to Baffin Bay**: North to northwest winds (5-25 kn, 3-13 m/s) today and Tuesday. South to southwest winds (5-15 kn, 3-8 m/s) Tuesday evening through Friday.

To see previous bulletins and forecasts for other Harmful Algal Bloom Bulletin regions, visit the NOAA Harmful Algal Bloom Operational Forecast System bulletin archive: http://tidesandcurrents.noaa.gov/hab/bulletins.html



Verified and suspected HAB areas shown in red. Other areas of high chlorophyll concentration shown in yellow (see p. 1 analysis for interpretation).